

AMENDMENTS TO THE CLAIMS

Claims 1-33 (Canceled)

34. (Currently Amended) An in vitro method for detection of disorders characterized by abnormal cell proliferation in an individual comprising

a. detecting in each of a biological test sample obtained from said individual and a normal control sample, a level of polynucleotides expression that hybridize under stringent conditions to probes specific for ~~of~~ a transketolase like-1 gene, wherein said probes whose complement hybridizes ~~hybridize~~ under stringent conditions to SEQ ID NO: 1; and

b. comparing said level of polynucleotides expression from said biological test sample to said level of polynucleotides expression in the normal control sample, wherein a higher level of polynucleotides expression in ~~the said~~ biological test sample as compared to said level of polynucleotides expression in ~~the said~~ normal control sample indicates that said individual has at least one of said disorders characterized by abnormal cell proliferation.

35. (Previously Presented) The method according to claim 34, wherein the disorder characterized by abnormal cell proliferation is cancer.

36. (Previously Presented) The method according to claim 35, wherein the cancer is colon cancer, lung cancer, gastric cancer or pancreatic cancer.

37. (Previously Presented) The method according to claim 34, wherein the biological test sample is a body fluid, a secretion, a smear, a biopsy, a liquid containing cells, lysed cells, cell debris, peptides or nucleic acids.

38. (Previously Presented) The method according to claim 37, wherein the biological test sample is serum, urine, semen, stool, bile, a biopsy or a cell- or tissue-sample.

39. (Withdrawn) The method according to claim 34, wherein the detection of the expression of the human transketolase like-1 gene is carried out on a polypeptide level.

40. (Canceled)

41. (Withdrawn) The method according to claim 39, wherein the detection on the polypeptide level is carried out using a binding agent directed against human transketolase like-1 polypeptides.
42. (Withdrawn) The method of claim 41, wherein the binding agent is an antibody, a fragment of an antibody, a peptidomimetic comprising an antigen binding epitope or a mini-antibody.
43. (Withdrawn) The method according to claim 39, wherein the detection is an immunocytochemical detection procedure.
44. (Previously Presented) The method according to claim 34, wherein at least one of steps (a) and (b) comprises using at least one nucleic acid probe, that hybridizes under stringent conditions to SEQ ID NO: 1.
45. (Previously Presented) The method according to claim 44, wherein the probe is detectably labeled.
46. (Previously Presented) The method according to claim 45, wherein the label is selected from the group consisting of a radioisotope, a bioluminescent compound, a chemiluminescent compound, a fluorescent compound, a metal chelate, and an enzyme.
47. (Previously Presented) The method according to claim 34, wherein at least one of steps (a) and (b) comprises using a nucleic acid amplification reaction.
48. (Previously Presented) The method according to claim 47, wherein the amplification reaction is selected from the group consisting of PCR, LCR and NASBA.
49. (Previously Presented) The method according to claim 44, wherein at least one of steps (a) and (b) comprises hybridizing the at least one nucleic acid probe in-situ.
50. (Previously Presented) The method according to claim 34, wherein at least one of steps (a) and (b) comprises performing in vitro molecular imaging.
51. (Withdrawn) A kit for performing the method of claim 34, which is a research kit or a diagnostic kit.

52. (Withdrawn) The kit of claim 51 comprising
- a. at least one probe for the detection of human transketolase like-1 gene expression products in biological samples;
 - b. a human transketolase like-1 gene product sample for performing a positive control reaction.
53. (Withdrawn) The kit of claim 52, wherein the probe is a nucleic acid probe, specifically hybridizing to human transketolase like-1 nucleic acids or an antibody specifically binding human transketolase like-1 proteins.
54. (Withdrawn) A method for treating disorders characterized by abnormal proliferation of cells based on the administration of a pharmaceutical composition containing a human transketolase like-1 gene or gene product in a pharmaceutical acceptable form.
55. (Withdrawn) The method according to claim 54, wherein the human transketolase like-1 gene or gene product is a nucleic acid in sense or antisense orientation or a polypeptide.
56. (Withdrawn) The method according to claim 55, wherein the pharmaceutical composition comprises a chimeric nucleic acid comprising a human transketolase like-1 nucleic acid or fragments thereof or a fusion polypeptide comprising a human transketolase like-1 polypeptide or fragments thereof.
57. (Withdrawn) The method according to claim 54, wherein the disorder characterized by abnormal cell proliferation is cancer.
58. (Withdrawn) The method according to claim 56, wherein the cancer is colon cancer, lung cancer, gastric cancer or pancreatic cancer.
59. (Withdrawn) The method according to claim 54, wherein the method for treatment is immunotherapy.
60. (Withdrawn) The method according to claim 54, wherein the method for treatment is vaccination therapy.

61. (Withdrawn) A method of identifying and obtaining a drug candidate for therapy of tumors of the colon, the lung, the pancreas or the stomach comprising the steps of
- a. contacting a TKT-L1 polypeptide as used in the method of the present invention or a cell expressing said polypeptide in the presence of components capable of providing a detectable signal in response to transketolase activity or to altered regulation of cell proliferation, and
 - b. detecting presence or absence of a signal or increase of the signal generated from transketolase activity or altered regulation of cell proliferation, wherein the absence or decrease of the signal is indicative for a putative drug.
62. (Withdrawn) A pharmaceutical composition for the treatment of tumors of the colon, the lung, the pancreas or the stomach, comprising a compound identifiable by the method according to claim 61, an antithiamine compound, an inhibitor of transketolase enzyme activity, an inhibitor of transketolase like-1 activity, a transketolase like-1 polypeptide or a human transketolase like-1 nucleic acid.
63. (Withdrawn) A method for rational tumor management comprising
- a. detecting the presence or absence and or the level of overexpression of transketolase like-1 gene in biological samples
 - b. building of subgroups according to the presence or absence and/or the levels of transketolase like-1 gene
 - c. tailoring an adequate therapy according to the subgroups comprising reduction of transketolase like-1 activity in individuals or in cells of individuals.
64. (Withdrawn) The method according to claim 63, wherein the reduction of the activity of transketolase like-1 is achieved by the administration of antithiamine compounds, of pharmaceutical compositions of claim 63, of inhibitors of transketolase enzyme activity, of transketolase like-1 antisense constructs, of ribozymes specific for transketolase like-1 or by reduced administration of thiamine.
65. (New) The method according to claim 34 wherein SEQ ID NO: 1 is said transketolase like-1

U.S. Application No.: 10/511,813
Attorney Docket: 4007-008

gene of which the level of polynucleotides is detected.

66. (New) The method according to claim 36 wherein SEQ ID NO: 1 is the transketolase like-1 gene of which the level of polynucleotides is detected.

67. (New) The method according to claim 36, wherein the cancer is colon cancer.

68. (New) The method according to claim 66, wherein the cancer is colon cancer.